



He's frequency formulation for the relativistic harmonic oscillator

Xu-Chu Cai^{a,*}, Wen-Ying Wu^b

^a College of Sciences, Donghua University, Shanghai 201620, China

^b College of Mechanical Engineering, Donghua University, Shanghai 201620, China

ARTICLE INFO

Keywords:

Relativistic harmonic oscillator
Periodic solution
He's frequency formulation

ABSTRACT

In this paper, He's frequency formulation is used to solve the nonlinear differential equation that governs the oscillation of the relativistic oscillator. The obtained solution with high accuracy shows that He's frequency formulation is a simple but promising method without any requirement for advanced calculus.

Crown Copyright © 2009 Published by Elsevier Ltd. All rights reserved.

Consider the following nonlinear differential equation of a relativistic oscillator [1]

$$\frac{d^2 u}{dt^2} + \frac{u}{\sqrt{1+u^2}} = 0 \quad (1)$$

with initial conditions

$$u(0) = B, \quad u'(0) = 0. \quad (2)$$

According to He's frequency formulation [2–5], we have

$$\omega^2 = \frac{\omega_1^2 R_2(t_2) - \omega_2^2 R_1(t_1)}{R_2(t_2) - R_1(t_1)} \quad (3)$$

where t_1 and t_2 are location points. Generally we set [6]

$$t_1 = \frac{T_1}{12}, \quad t_2 = \frac{T_2}{12} \quad (4)$$

where T_1 and T_2 are periods of the trial solutions $u_1(t) = B \cos t$ and $u_2(t) = B \cos \omega t$, respectively, which lead to the following residuals:

$$R_1(t_1) = -B \cos t_1 + \frac{B \cos t_1}{\sqrt{1+B^2 \cos^2 t_1}}, \quad (5)$$

$$R_2(t_2) = -B \omega^2 \cos \omega t_2 + \frac{B \cos \omega t_2}{\sqrt{1+B^2 \cos^2 \omega t_2}}. \quad (6)$$

Using Eq. (3), we obtain:

$$\omega^2 = \frac{\omega_1^2 R_2(t_2) - \omega_2^2 R_1(t_1)}{R_2(t_2) - R_1(t_1)} = \frac{-B \omega^2 \cos \frac{\omega T_2}{12} + \frac{B \cos \frac{\omega T_2}{12}}{\sqrt{1+B^2 \cos^2 \frac{\omega T_2}{12}}} - \omega^2 \left(-B \cos \frac{T_1}{12} + \frac{B \cos \frac{T_1}{12}}{\sqrt{1+B^2 \cos^2 \frac{T_1}{12}}} \right)}{-B \omega^2 \cos \frac{\omega T_2}{12} + \frac{B \cos \frac{\omega T_2}{12}}{\sqrt{1+B^2 \cos^2 \frac{\omega T_2}{12}}} - \left(-B \cos \frac{T_1}{12} + \frac{B \cos \frac{T_1}{12}}{\sqrt{1+B^2 \cos^2 \frac{T_1}{12}}} \right)} = \frac{1}{\sqrt{1+\frac{3}{4}B^2}} \quad (7)$$

* Corresponding author.

E-mail addresses: caixuchu@dhu.edu.cn (X.-C. Cai), wywu@dhu.edu.cn (W.-Y. Wu).

which is the same as the first analytical approximate frequency obtained by Beléndez [1].

Hereby He's frequency formulation is proved to be a powerful mathematical tool in the search for periodic solutions of nonlinear oscillators without the requirement of perturbation or linearization; it is simple, straightforward and effective.

References

- [1] A. Beléndez, C. Pascual, A. Márquez, D.I. Méndez, Application of He's homotopy perturbation method to the relativistic (an) harmonic oscillator. I: Comparison between approximate and exact frequencies, *International Journal of Nonlinear Science and Numerical Simulation* 8 (4) (2007) 483–491.
- [2] J.-H. He, Some asymptotic methods for strongly nonlinear equations, *International Journal of Modern Physics B* 20 (10) (2006) 1141–1199.
- [3] J.-H. He, *Nonperturbative methods for strongly nonlinear problems*, Dissertation.de-Verlag im Internet GmbH. 2006.
- [4] J.-H. He, Comment on 'He's frequency formulation for nonlinear oscillators', *European Journal of Physics* 29 (2008) L1–L4.
- [5] J.-H. He, An improved amplitude-frequency formulation for nonlinear oscillators, *International Journal of Nonlinear Sciences and Numerical Simulation* 9 (2) (2008) 211–212.
- [6] Lei Geng, Xu-Chu Cai, He's frequency formulation for nonlinear oscillators, *European Journal of Physics* 28 (2007) 923–931.